



## FEATURES OF CHEMILUMINESCENCE OF NUTRIENTS DURING THE CULTIVATION OF BACTERIA

Nozimov Jakhongir Ikromjonovich

*Assistant, Namangan Institute of Engineering and Technology, Namangan region, Namangan city, st.  
Kasansay, Uzbekistan*

**Annotation:** *Bacteria of different species, due to the characteristics of their metabolism, have different abilities to counteract the formation of free radicals and oxidative stress. An indicator reflecting the development of oxidative processes is the antioxidant activity of nutrient media during bacterial cultivation.*

**Key words:** *microorganisms, chemiluminescence method, oxidative stress, free radicals, nutrient media*

### INTRODUCTION

*Pseudomonas aeruginosa* And *Escherichia coli* are gram-negative bacteria that are one of the most common pathogens of infectious diseases and have different structural, functional and metabolic characteristics. Some studies note different sensitivity of these microorganisms to antibiotics. For example, in the work of Hanberger H. et al . it has been shown that the sensitivity of *E. coli* to commonly used antibiotics remains high, while the susceptibility of *P. aeruginosa* is quite low [1]. In a study by Eagye KJ et al . low resistance of *E. coli* to antibiotics was noted compared to *P. aeruginosa* (resistance to imipenem 22–33%, to ciprofloxacin 5–21%) [2]. Along with the direct mechanisms of action of antibiotics, a mechanism that enhances their biocidal properties is the activation of oxidative stress in the bactericidal cell [3], caused by excessive formation of free radicals [4, 5]. Oxidative stress is a universal element of damage to molecules and cell structures, in particular, it has a destructive effect on the structure and activity of proteins, and also causes mutagenesis and death of bacterial cells [6, 7, 8, 9, 10, 11]. Bacterial cells have antioxidant systems to protect against oxidative stress. The higher the level of activity of these systems, the higher the resistance of microorganisms to oxidative stress. Assessing the potential to counteract oxidative processes may help explain the resistance of bacteria to antibacterial factors.

To maintain the vital activity of microorganisms, their normal growth and development, appropriate conditions are required. During cultivation, the most important nutrients and growth factors enter the bacterial cell as part of nutrient media (MS). PS are important for setting up experiments in various areas of scientific knowledge about microorganisms, and also serve as a necessary tool for solving clinical problems; they find application in clinical and laboratory diagnostics, microbiology, biotechnology, cell technologies, and bioengineering for tissue growth [12]. One of the parameters of the state of nutrient media is their antioxidant activity—the ability to counteract the formation of free radicals: reactive oxygen species (ROS) and lipid peroxide radicals [13]. A decrease in antioxidant activity in the culture medium may indicate the development of oxidative



stress in it, a decrease in the activity of protective enzyme systems of cells, and the active production of oxidative metabolites.

Among the methods for determining the antioxidant activity of substrates, one of the most sensitive is the method of recording chemiluminescence. Chemiluminescence (CL) is a luminescence that occurs during the interaction of free radicals, which can be selectively enhanced by the addition of various substances, in particular lucinogen and luminol [14].

#### Purpose of the study

Comparative assessment of the antioxidant activity of culture media samples during the cultivation of *P. aeruginosa* bacteria and *E. coli*.

The objects of the study were strains of the bacterium *P. aeruginosa* and *E. coli*, provided by the Clinic of the Federal State Budgetary Educational Institution of Higher Education BSMU of the Ministry of Health of Russia, isolated from a clinical sample. GRM broth was used as a nutrient medium. Standardization of the number of bacteria in the suspension was carried out by assessing the turbidity of the McFarland standard with an optical density of 1.0. The registration of chemiluminescence of nutrient media was assessed using a Chemiluminomer-003 device [14]. The method makes it possible to evaluate oxidative processes in various substrates, including cultivation media. The measurement principle is based on the detection of light quanta released during the interaction of extremely active agents - free radicals. The model in which the antioxidant activity of the culture media was assessed was a phosphate buffer ( $\text{KH}_2\text{PO}_4$  – 20 mM,  $\text{KCl}$  – 105 mM, pH 7.45 units), with the addition of sodium citrate (50 mM) and the phosphor luminol (5-amino-2,3-dihydro-1,4-phthalazinedione) to enhance the released light quanta. The CL assessment of the studied samples was carried out on days 1, 5 and 7. Samples of culture media in a volume of 0.5 ml were added to 20 ml of the model system and placed in the device chamber. After this, through a special hole in the device, a solution of iron sulfate ( $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  – 50 mM) was introduced into the mixture of media samples and the model system to initiate chain reactions of the formation of free radicals, primarily reactive oxygen species [15]. The samples were measured for 3 minutes and described in the CL kinetics parameters. To study not only the spontaneous antioxidant activity of the media, but also their state under conditions of oxidative stress, NaCl (50 mg/ml) was used. NaCl causes hyperosmotic stress and is used to hyperactivate oxidative processes. The CL parameters were expressed in conventional units of the integral index of the cross-sectional light sum. Statistical processing of the results was carried out using the Statistica for Windows software package. The normality of the distribution of actual data was checked using the Shapiro-Wilk test. Since the distribution differed from normal, the nonparametric Mann-Whitney test was calculated. The results were considered significant at  $p < 0.05$ . The indicators were presented as median (Me) and interquartile range (IQR).



REFERENCES:

- 1) Юсупова М. Н., Ахмедова М. М. МЕВАЛИ ДАРАХТЛАРНИ ЗАРАРКУНАНДАЛАРИГА УЙЎУНЛАШГАН КУРАШ ЧОРАЛАРИ //ЖУРНАЛ АГРО ПРОЦЕССИНГ. – 2020. – Т. 2. – №. 8.
- 2) Ходжаев, Ш. Т., Сагдуллаев, А. У., Исаев, О. Б., & Юсупова, М. Н. (2011). Проблемы защиты растений в Узбекистане. Защита и карантин растений, (8), 23-24. Юсупова М. Особенности защиты хлопчатника посеянного под пленки от вредных организмов //Автореф. канд. дисс./М. Юсупова–Ташкент. – 2001.
- 3) Ходжаев, Ш. Т., Юсупова, М. Н., Курязов, Ш., & Сагтаров, Н. (2008). Перспективы биологической защиты хлопчатника от хлопковой совки. Сб. трудов.- Ташкент: Таллин, 44-49.
- 4) Yusupova M. N., Nosirov B. Z. Pests of cotton and straw control at collection //EPRA International Journal of Multidisciplinary Research (IJMR)-Peer Reviewed Journal. – 2020. – Т. 6. – №. 12. – С. 57-61.
- 5) Yusupova M. N., Axmedova M. M. Mevali daraxtlarni zararkunandalariga uygunlashgan kurash choralari //Jurnal JURNAL AGRO PROTSESSING. Data publikatsii. – 2020. – №. 8. – С. 12.
- 6) Yusupova M. N. Biological method of crop protection in the fergana valley //Agrarian science. – 2018. – №. 6. – С. 68-70.
- 7) Urmonovich, Numonov Otabek. "MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES." ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ 14.5 (2023): 3-5.
- 8) MN, Yusupova, and B. Z. Nosirov. "Control Of Cotton Pests On Stubble Lands." International Journal of Applied 10.2 (2015): 99-108.
- 9) Юсупова М. Н., Тургунова А. Н., Очилов С. Н. Система интегрированной защиты растений //Российский электронный научный журнал. – 2015. – №. 1. – С. 169-174.
- 10) Alimzhanova Z. I., Kadyrova D. S., Yusupova M. N. Ceramic pigments based on raw materials from Uzbekistan //Glass and Ceramics. – 2014. – Т. 70. – №. 11-12. – С. 441-443.
- 11) Yusupova M. N., Gapparov A. M. Biological Method Of Plant Protection In Uzbekistan //The American Journal of Agriculture and Biomedical Engineering. – 2020. – Т. 2. – №. 11. – С. 29-32.
- 12) Rashidovna M. N., Urmonovich N. O. Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types //International Journal on Orange Technologies. – Т. 2. – №. 10. – С. 79-82.
- 13) Юсупова М. Н., Носиров Б. З. БИОЛОГИЧЕСКИЙ МЕТОД ЗАЩИТЫ РАСТЕНИЙ В УЗБЕКИСТАНЕ //Научно-практические пути повышения экологической устойчивости и социально-экономическое обеспечение сельскохозяйственного производства. – 2017. – С. 498-501.



- 14) Urmonovich, N. O. (2023). MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES. ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ, 14(5), 3-5.
- 15) Yusupova M. et al. Protection of after harvest cultures-as a reservetors of cotton pests //Agriculture and Biology Journal of North America. – 2013. – Т. 4. – №. 5. – С. 576-582.
- 16) Ходжаев, Ш. Т., Юсупова, М. Н., Юлдашев, Ф., Исаев, О. Б., & Шокирова, Г. (2011). Борьба с вредителями хлопчатника на пожнивных культурах в севообороте. Вестник защиты растений, (2), 46-52.
- 17) Yusupova M. N. et al. Possibilities of the biological method of cotton plant protection //Agriculture and Biology Journal of North America. – 2011. – Т. 2. – №. 5. – С. 742-744.
- 18) Ходжаев, Ш. Т., Юсупова, М. Н., Юлдашев, Ф., & Жамалов, А. Г. (2010). Хлопковая совка на пожнивных культурах. Защита и карантин растений, (12), 22-23.
- 19) Хайдарова, Х. А., Юсупова, М. Н., Ихтиярова, Г. А., & Хайдаров, А. А. ПОЛУЧЕНИЕ ХИТОЗАНА ИЗ ПОДМОРА ПЧЕЛ APIS MILLIFERA. Сучасний рух науки: тези доп. XI міжнародної науково-практичної інтернет-конференції, 8-9 жовтня 2020 р.-Дніпро, 2020.-Т. 2.-426 с., 352.
- 20) Yusupova M., Turgunova A., Ochilov S. INTERGRATED PLANT PROTECTION SYSTEMS.
- 21) Abduhamidovich N. A. et al. MANGOSTIN DARAXTI VA MEVASINI TIBBIYOTDA FOYDALANISH //Journal of new century innovations. – 2023. – Т. 28. – №. 2. – С. 12-14.
- 22) Urmonovich N. O. MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES //ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ. – 2023. – Т. 14. – №. 5. – С. 3-5.
- 23) Юсупова, Махпуза Нумановна. "АНОРНИ ЗАРАРКУНАНДАЛАРДАН ҲИМОЯЛАШ." PEDAGOG 6.4 (2023): 562-567.
- 24) Юсупова М. Н. и др. ФАРҒОНА ВОДИЙСИ ШАРОИТИДА ИГНА БАРГЛИ ДАРАХТЛАРНИ ЗАРАРКУНАНДАЛАРДАН ҲИМОЯЛАШ //SO 'NGI ILMİY TADQIQOTLAR NAZARIYASI. – 2023. – Т. 6. – №. 4. – С. 316-320.